



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Molecular characterization of blaESBL-producing *Escherichia coli* cultured from pig farms in Ireland

Wang, J ; Gibbons, James F ; McGrath, K ; Bai, L ; Li, Fengqin ; Leonard, Finola C ; Stephan, Roger ; Fanning, Séamus

Abstract: Objectives: To characterize ESBL-encoding *Escherichia coli* cultured from pigs and their plasmids carrying these genes following conjugation into recipient strains. **METHODS:** Six ESBL-producing *E. coli* were recovered from faecal samples taken from pigs along with a further isolate from the environment of a farrowing house on three pig farms in Ireland. These isolates were characterized by phylogenetic grouping, MLST and ESBL genotype analyses. Conjugation experiments were carried out in broth mating assays. S1-nuclease PFGE was used to determine the plasmid profiles. Whole-genome sequences of the seven *E. coli* were determined and subsequently analysed. **RESULTS:** Phylogenetic groups and the corresponding MLST STs identified among the seven tested *E. coli* isolates included A/ST10, A/ST34, C/ST23 and C/ST1629. All seven isolates carried one or more high-molecular-weight plasmids and demonstrated the ability to transfer their cefotaxime resistance phenotype at high frequencies. Five transmissible plasmid replicon types were detected, including IncK/B (n = 3), IncI1 (n = 2), IncFIA (n = 1), IncFIB (n = 1) and IncN (n = 1). ESBL-encoding genes, including blaCTX-M-14, blaCTX-M-15 and blaTEM-20, were identified. **CONCLUSIONS:** As the first report from pig sources in Ireland, characterization of these ESBL-encoding isolates and their transmissible plasmids extends our understanding on these resistance markers from porcine *E. coli*.

DOI: <https://doi.org/10.1093/jac/dkw278>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-130012>

Journal Article

Accepted Version

Originally published at:

Wang, J; Gibbons, James F; McGrath, K; Bai, L; Li, Fengqin; Leonard, Finola C; Stephan, Roger; Fanning, Séamus (2016). Molecular characterization of blaESBL-producing *Escherichia coli* cultured from pig farms in Ireland. *Journal of Antimicrobial Chemotherapy*, 71(11):3062-3065.

DOI: <https://doi.org/10.1093/jac/dkw278>

JAC-2016-0189 revision 1, as a brief note-

Molecular characterization of *bla*_{ESBL}-producing *Escherichia coli* cultured from pig farms in Ireland

Juan Wang^{1,2}, James F. Gibbons³, Kathleen McGrath², Li Bai⁴, Fengqin Li⁴, Finola C. Leonard³, Roger Stephan⁵ and Séamus Fanning^{2,4,6}

¹College of Veterinary Medicine, Northwest A&F University, Yangling 712100, Shaanxi, China;

²UCD-Centre for Food Safety, School of Public Health, Physiotherapy & Sports Science, University College Dublin, Belfield, Dublin 4, Ireland;

³School of Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland;

⁴Key Laboratory of Food Safety Risk Assessment, Ministry of Health, China National Centre for Food Safety Risk Assessment, 7 Panjiayuan Nanli, Chaoyang District, Beijing 100021, China;

⁵Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 272, CH-8057 Zurich, Switzerland;

⁶Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, Stranmillis Road, Belfast BT9 5AG, Northern Ireland

Corresponding author- Tel.: +353-1-716 2869; Fax: +353-1-716 1147; sfanning@ucd.ie

37 **Sir,**

38 ESBL-producing *Escherichia coli* of porcine origin have been detected with
39 increasing frequency during the past few years, and plasmid-mediated transfer
40 of ESBL genes from pigs is considered to be a significant factor driving the
41 dissemination of these markers.^{1, 2} However, epidemiological data describing
42 ESBL-producing *E. coli* in Irish food-producing animals is limited. In an earlier
43 study, the antimicrobial resistance associated with a total of 1,126 *E. coli*
44 cultured from 39 Irish pig farms, located in geographically distinct regions, with
45 recorded variability in duration of in-feed antimicrobial compound use, was
46 studied.³ This current report describes a follow-up study, in which seven
47 isolates that were resistant to cefotaxime were further characterized.

48
49 The corresponding serotypes; phlotypes and sequence types were
50 determined for all seven isolates studied, using previously published protocols
51 (**Table 1**).^{4, 5} The MIC for cefotaxime was determined by Etest and
52 ESBL-encoding genes were further characterized by PCR and amplicon
53 sequencing.⁶ Conjugation experiments were carried out on each isolate
54 individually to assess their ability to transfer cefotaxime resistance.⁷
55 Antimicrobial susceptibility testing was performed on the *E. coli* and their
56 associated transconjugants by agar disc diffusion according to the criteria of
57 CLSI.⁸ Plasmid analysis included replicon typing and S1-nuclease PFGE
58 molecular profiling methods.⁷ WGS was undertaken for all seven isolates
59 using the Illumina HiSeq 2500 platform. The core-genome based phylogeny
60 was determined by Harvest Suite and a maximum composite likelihood
61 phylogenetic tree was generated with MEGA version 6 software using *E. coli*
62 MG1655 as reference.^{9, 10} All genomic data was submitted to EMBL and
63 Accession numbers are listed in **Table 1**.

64
65 The source origins of all seven ESBL-positive *E. coli* are shown in **Table 1** and
66 these were composed of four different serotypes that could be assigned into
67 two phylogenetic groups and four sequence types. Four of the seven isolates
68 were identified as ST10 and ST34, belonging to the CC10, which is one of the
69 largest clonal complexes identified within the *E. coli* MLST database. These
70 isolates were cultured from two farms that were geographically located in
71 separate regions, Cork and Roscommon. The latter ESBL-positive *E. coli*
72 possessed *bla*_{CTX-M-15} and *bla*_{TEM-20} –encoding genes (**Table 1**). The important
73 contribution of the ST10 type to the spread of resistance is a very recent
74 observation; wherein previously this sequence type was reported both in
75 humans and/or animals in Spain, Italy, Germany and other European
76 countries.¹¹⁻¹³ Similarly, members of the ST23 clonal complex have also been
77 observed to be an important source of *bla*_{CTX-M} genes.¹³ In our study, two ST23
78 isolates were identified from a single farm and carried *bla*_{CTX-M-14}. The same
79 ESBL-genotype was also linked with a ST1629 from a separate group of
80 animals on the same farm in this study. To the best of our knowledge, this is

the first reported identification of ST1629 carrying ESBL genes from food-producing animals.

All isolates transferred their cefotaxime resistance marker to a susceptible *E. coli* 26R 793 (Rif^r) recipient with transfer rates ranging from 3.2×10^{-4} to 8.5×10^{-1} . The MIC determined for cefotaxime by Etest recorded values ranging from 8- to >32- mg/L. The *bla*_{ESBL} genes transferred *via* conjugation were subsequently confirmed as *bla*_{TEM-1}, *bla*_{TEM-20}, *bla*_{CTX-M-14}, and *bla*_{CTX-M-15} (Table 1). The first study to report on the prevalence of ESBL-positive Enterobacteriaceae in Ireland, was conducted in 2003, but was limited to clinical settings.¹⁴ These data showed that *bla*_{CTX-M-15} and *bla*_{CTX-M-14} genes were the most common ESBL genotypes identified in those bacterial isolates cultured from Irish hospitals.¹⁵ The same genotypes were also recovered in Ireland from *Salmonella enterica* from humans.¹⁶ A *bla*_{TEM-20} genotype was identified in our study and this was previously identified in a *Salmonella enterica* (serovar Java) cultured from Irish beef in 2006.¹⁶

Plasmid profiles were determined by S1-nuclease treatment followed by PFGE. All isolates contained detectable large molecular-weight plasmids.

Heterogeneity among the profiles was a common feature noted, although most of the plasmid profiles were related among strains isolated from same geographical locations (Figure S1). Following conjugation, the plasmid profiles were again determined by S1-nuclease PFGE for the transconjugants recovered. Two isolates from Cavan, F1B2 and F1L4 possessed one transmissible plasmid (of approximately 100-kbp). Two plasmids were transferred in the case of isolate F1L3, and which were approximately 120- and 200-kbp in size. From the isolates F13P5 and F13P4, only one transmissible plasmid was detected in each case, of between 70- and 80-kbp, respectively. The remaining two isolates, F25OS1 and F25P5, possessed two large transmissible plasmids. Three replicon types (including IncFIA, IncFIB and IncK/B) were detected in *E. coli* F1L3 and all three Inc-types were also identified from the corresponding transconjugant. The same Inc-type profile was also noted in *E. coli* F1L4, and IncK/B was the sole type detected following conjugation (Table 1). In the *E. coli* isolates denoted as F25OS1 and its transconjugant, IncI1 and IncN types were identified. Bacterial isolates, F13P5 and F13P4 that were originally cultured from piglets in Cork, were positive for an IncFIB type plasmid. The IncK/B and IncI1 types were detected in conjugative mating pairs for F1B2 and F25P5, respectively (Table 1).

In an earlier report, conjugative pCT-like plasmids (of the IncK group, approximately 100-kbp in size, accession no. FN868832) were determined to be important vectors contributing to the dissemination of *bla*_{CTX-M-14} genes in the United Kingdom, Europe, Australia, and Asia.^{17, 18} In our study, three isolates carried these *bla*_{CTX-M-14} genes in association with IncK/B type

plasmids. Of these, one transmissible plasmid of approximately 100-kbp was also identified. Furthermore, according to the WGS data, 78%, 78% and 95% of the plasmid genetic backbone of pCT were also present in all three *bla*_{CTX-M-14}-producing isolates from a single farm, namely F1L3, F1L4 and F1B2, respectively. Based on these observations, it is tempting to speculate that pCT-like plasmids may be contributing to the spread of *bla*_{CTX-M-14} genes among these pig herds in Ireland. Further sequence analysis and reconstruction of these plasmids would be required to support this conclusion, a feature that will be the subject of a separate report.

In order to assess the genetic relationships between these ESBL-producing *E. coli* isolates, a phylogenetic tree based on their core-genome was generated with MEGA6 software. **Figure-1** shows that the seven ESBL strains were clustered into three different groups. The seven *E. coli* clustered according to their ST. Four ST10 complex (including ST34) *E. coli*, which were cultured from two locations in Ireland, formed a closely related cluster with *E. coli* MG1655. All of these isolates were determined to be phylogroup A isolates. *E. coli* of sequence type ST1629 are rarely reported, and this isolate was separated by a large genetic distance when compared with the other two clusters identified. Furthermore, analysis of the genomic data also demonstrated that these *E. coli* shared several common virulence factors, such as *fimFGH* (type I fimbriae, Adherence), *ibeBC* (CFA/I fimbriae, Adherence), *ehaB* (AIDA-I type, Autotransporter), *espL/X* (Non-locus-of-enterocyte-effacement encoded type-three-secretion-system effectors) and *hlyE/clyA* (hemolysin/cytolysin A, Toxins) [**Table S1**, available as Supplementary data at JAC Online].¹⁹

In conclusion, our findings documented the characterization and potential for dissemination of their *bla*_{CTX-M}-encoding genes from *E. coli* cultured from pig farms in Ireland. The ST10 complex *E. coli* from two different locations, appeared to demonstrate an association with *bla*_{CTX-M-15} and *bla*_{TEM-20} genes. Isolates of ST23 and ST1629 sequence types from the same location were associated with a *bla*_{CTX-M-14} gene. All ESBL-encoding genes studied could be transferred to a recipient bacterium *via* conjugation. WGS data showed that bacterial isolates belonging to the ST10 complex in this collection were closely related to *E. coli* MG1655, and that pCT-like plasmids identified may contribute to the transmission of *bla*_{CTX-M-14} genes. Further studies are required to evaluate potential zoonotic links between ESBL-producing *E. coli* from pigs and human cases of infection.

Funding-

This study was supported by internal funding.

Transparency declarations-

None to declare.

References-

1. Hammerum AM, Larsen J, Andersen VD *et al.* Characterization of extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* obtained from Danish pigs, pig farmers and their families from farms with high or no consumption of third- or fourth-generation cephalosporins. *J Antimicrob Chemother* 2014; **69**: 2650-7.
2. Geser N, Stephan R, Kuhnert P *et al.* Fecal carriage of extended-spectrum beta-lactamase-producing Enterobacteriaceae in swine and cattle at slaughter in Switzerland. *J Food Prot* 2011; **74**: 446-9.
3. Gibbons JF, Boland F, Egan J *et al.* Antimicrobial resistance of faecal *Escherichia coli* isolates from pig farms with different durations of in-feed antimicrobial use. *Zoonoses public health* 2015; **63**: 241-50.
4. Wirth T, Falush D, Lan R *et al.* Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* 2006; **60**: 1136-51.
5. Clermont O, Christenson JK, Denamur E *et al.* The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep* 2013; **5**: 58-65.
6. Dallenne C, Da Costa A, Decre D *et al.* Development of a set of multiplex PCR assays for the detection of genes encoding important beta-lactamases in Enterobacteriaceae. *J Antimicrob Chemother* 2010; **65**: 490-5.
7. Wang J, Stephan R, Karczmarczyk M *et al.* Molecular characterisation of *bla*_{ESBL}-harbouring conjugative plasmids identified in multi-drug resistant *Escherichia coli* isolated from food-producing animals and healthy humans. *Front Microbiol* 2013; **4**: 188.
8. CLSI. Performance standards for antimicrobial susceptibility testing; Twenty-fourth informational supplement M100-S24. 2014, Wayne, PA: Clinical and Laboratory Standards Institute.
9. Tamura K, Stecher G, Peterson D *et al.* MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 2013; **30**: 2725-9.
10. Treangen TJ, Ondov BD, Koren S *et al.* The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol* 2014; **15**: 524.
11. Izdebski R, Baraniak A, Fielt J *et al.* Clonal structure, extended-spectrum beta-lactamases, and acquired AmpC-type cephalosporinases of *Escherichia coli* populations colonizing patients in rehabilitation centers in four countries. *Antimicrob Agents Chemother* 2013; **57**: 309-16.
12. Oteo J, Diestra K, Juan C *et al.* Extended-spectrum beta-lactamase-producing *Escherichia coli* in Spain belong to a large variety of multilocus sequence typing types, including ST10 complex/A, ST23 complex/A and ST131/B2. *Int J Antimicrob Agents* 2009; **34**: 173-6.

- 212 13. Naseer U, Sundsfjord A. The CTX-M conundrum: dissemination of plasmids
213 and *Escherichia coli* clones. *Microb Drug Resist* 2011; **17**: 83-97.
- 214 14. Morris D, O'Hare C, Glennon M *et al*. Extended-spectrum beta-lactamases in
215 Ireland, including a novel enzyme, TEM-102. *Antimicrob Agents Chemother* 2003;
216 **47**: 2572-8.
- 217 15. Walsh F, Rogers TR. Comparison of plasmid-mediated quinolone resistance
218 and extended-spectrum beta-lactamases in third-generation
219 cephalosporin-resistant Enterobacteriaceae from four Irish hospitals. *J Med*
220 *Microbiol* 2012; **61**: 142-7.
- 221 16. Morris D, Whelan M, Corbett-Feeney G *et al*. First report of
222 extended-spectrum-beta-lactamase-producing *Salmonella enterica* isolates in
223 Ireland. *Antimicrob Agents Chemother* 2006; **50**: 1608-9.
- 224 17. Stokes MO, Cottell JL, Piddock LJV *et al*. Detection and characterization of
225 pCT-like plasmid vectors for *bla*_{CTX-M-14} in *Escherichia coli* isolates from humans,
226 turkeys and cattle in England and Wales. *J Antimicrob Chemoth* 2012; **67**:
227 1639-44.
- 228 18. Cottell JL, Webber MA, Coldham NG *et al*. Complete sequence and molecular
229 epidemiology of IncK epidemic plasmid encoding *bla*_{CTX-M-14}. *Emerg Infect Dis*
230 2011; **17**: 645-52.
- 231 19. Chen L, Zheng D, Liu B *et al*. VFDB 2016: hierarchical and refined dataset for
232 big data analysis-10 years on. *Nucleic Acids Res* 2016; **44**: D694-7.